## Colonoscopy Miralax Prep Instructions AM

## DATE/TIME OF PROCEDURE – PLEASE CHECK THE EMAIL THESE INSTRUCTIONS WERE ATTACHED TO

\*\*Your Arrival time is subject to change. The Surgery Center will contact you approximately 2 days before your procedure to confirm Arrival Time\*\*

LOCATION: East Bay Endosurgery Center

300 Frank H. Ogawa Plaza, Suite 135, Oakland, CA 94621

Phone: (510) 893-1600

LATE CANCELLATION (LESS THAN 3 DAYS) FEE OF \$200 – If you need to cancel, please call (510) 444-3297

What to Do

prior to Procedure

What to do 5 days

What to do the day before your procedure On The Day of the Procedure



You will need to buy the following medications:

7 days Prior to Procedure

- Miralax 510 grams (17.9 oz) powder or generic polyethylene glycol 3350 (can find these in the laxative section of the store.

AND

- One (1) gallon of water, Gatorade, Propel, Crystal Light or other noncarbonated clear liquid (no red color)

AND

- Four (4) Simethicone tablets (i.e. Gas-Ex)
- Four (4) Bisacodyl 5 mg tablets
- If you are diabetic, you may use sugar-free choices

Review your prep instructions thoroughly.

Please continue all medications including aspirin. If you are taking a blood thinner, please contact the prescribing physician for instructions.



STOP eating any raw vegetables or vegetables containing seeds, fruit with seeds or skin, corn, popcorn, nuts, quinoa, beans, and seeds.



Please arrange a driver to take you home. This needs to be family or a friend. Uber, Lyft, and taxi are only allowed if you have a family or friend accompany you. You will not receive sedation without a family or friend available to drive vou home. All rides must be available by 3:30pm.



If you are taking insulin for diabetes, you will receive separate instructions.

## NO SOLID FOODS

Clear Liquids only all day. 9am – Prepare one (1) gallon of preferred clear liquid with 16 capfuls of Miralax. Stir until dissolved. Chill if desired. 2pm – Drink first half of the Miralax solution (64 ounces or 8 cups). Drink one (1) 8-ounce glass every 15 minutes until complete.

Take two (2) 5 mg Bisacodyl tablets and two (2) 125 mg Simethicone tablets (i.e.-Gas Ex) followed by (2) 8 oz. glasses of water.

**7pm** – Drink the second half of the Miralax-Gatorade solution. Drink one (1) 8-ounce glass every 15 minutes until completed.

Take two (2) 5 mg of Bisacodyl and two (2) 125 mg Simethicone tablets (i.e.-Gas Ex) followed by (2) 8 oz. glasses of water.

After you are finished with the solution, continue clear liquids only.

Drink plenty of water and clear liquids throughout the day to avoid dehydration.

Nothing by mouth starting two (2) hours prior to your Arrival Time. No Gum or hard candy.

If you take daily medication, please take with a small sip of water at least two (2) hours before your arrival time, unless otherwise instructed.



## Reminders

- If you use an inhaler, bring it with you to your procedure
- Please do not wear any jewelry and remove piercings
- Bring insurance cards
- Be prepared to pay your copay

Your ride will not be able to stay here after you are dropped off, but we will need a phone number for them to pick you up after your procedure. Most patients are ready for pick up approximately two (2) hours after dropping off; however, medical procedure timing is unpredictable, and this may vary. Please do not make any timesensitive commitments shortly after your procedure.

If you have had any procedures, started any new medications, or had a change in your medical status since scheduling this procedure, please contact our office at (510) 444-3297

\*Clear Liquid Diet Details: NO RED COLOR. NO DAIRY

Approved: Clear sodas, black coffee, tea, clear juices, clear fitness waters, chicken broth, vegetable broth, beef broth, Gelatin, Popsicles without pulp, sorbet without pulp







**Avoid** No milk/dairy No juices with pulp Popsicles with pulp **NO RED COLOR**.

