Colonoscopy TRILYTE (GAVILYTE, GOLYTELY) Split Dose Prep Instructions

DATE/TIME OF PROCEDURE – PLEASE CHECK THE EMAIL THESE INSTRUCTIONS WERE ATTACHED TO

Your Arrival time is subject to change. The Surgery Center will contact you approximately 2 days before your procedure to confirm Arrival Time

LOCATION: East Bay Endosurgery Center

300 Frank H. Ogawa Plaza, Suite 135, Oakland, CA 94621

Phone: (510) 893-1600

LATE CANCELLATION (LESS THAN 3 DAYS) FEE OF \$200 – If you need to cancel, please call (510) 444-3297

What to do 7 days Prior to Procedure

What to do 5 days prior to Procedure

What to do the day before your procedure

On the Day of the procedure



Please make sure you have picked up the TriLyte (or GaviLyte, or GoLytely) prep medication from your pharmacy noted above.

Review your prep instructions thoroughly.

Please continue all medications including aspirin.

If you are taking a blood thinner, please contact the prescribing physician for instructions.



Please arrange

a driver to take you home. This needs to be a family or friend. Uber, Lyft, and taxi are only allowed if you have a family or friend accompany you. You will not receive sedation without a family or friend available to drive you home. All rides must be available by 3:30pm.



STOP eating any raw vegetables or vegetables containing seeds, fruit with seeds or skin, corn, popcorn, nuts, quinoa, beans and seeds.

Purchase clear liquid diet items. (See page 2 for clear liquid items)

Please Purchase the following over the counter medication:

1. (4) 125 mg Simethicone tablets (i.e.-Gas Ex)



If you are

taking insulin for diabetes, you will receive separate instructions



NO SOLID FOODS

Clear Liquids only upon waking up in the morning. A clear liquid diet is necessary for a colonoscopy.

Prep:

5pm-Please drink ⅓ of your prep solution (64 oz)

After the prep is complete take (2) 125 mg Simethicone tablets (i.e.-Gas Ex) followed by (2) 8oz glasses of water.

Drink plenty of water and clear liquids throughout the day to avoid dehydration.

Continue a clear liquid diet.

Drink the remaining ½ of your prep solution (64 oz) four (4) hours before your Arrival Time.

After completing prep take (2) 125 mg Simethicone tablets (i.e.-Gas Ex) followed by (2) 8 oz. glasses of water.

Nothing by mouth starting two (2) hours prior to your Arriva Time. No Gum or hard candy.

If you take daily medication, please take with a small sip of water at least two (2) hours prior to Arrival Time, unless otherwise instructed.



Reminders

- If you use an inhaler, bring it with you to your procedure
- Please do not wear any jewelry and remove piercings
- Bring insurance cards
- Be prepared to pay your copay

Your ride will not be able to stay here after you are dropped off, but we will need a phone number for them to pick you up after your procedure. Most patients are ready for pick up approximately two (2) hours after dropping off; however, medical procedure timing is unpredictable, and this may vary. Please do not make any time-sensitive commitments shortly after your procedure.

If you have had any procedures, started any new medications, or had a change in your medical status since scheduling this procedure, please contact our office at (510) 444-3297

*Clear Liquid Diet Details: NO RED COLOR LIQUIDS. NO DAIRY

Approved: Clear sodas, black coffee, tea, clear juices, clear fitness waters, Ensure Clear, chicken broth, vegetable broth, beef broth, Gelatin, Popsicles without pulp, sorbet without pulp



Avoid No milk/dairy No juices with pulp Popsicles with pulp **NO RED COLOR.**

